

## Venous Thromboembolism Prophylaxis in Surgical Patients

(2024-2027)

### REFERENCES

1. Gould MK, Garcia DA, Wren SM, et al. Prevention of VTE in nonorthopedic Surgical Patients. *Chest*. 2012;141(2 Suppl):e227S-e277S. doi:10.1378/chest.11-2297
2. Anderson DR, Morgano GP, Bennett C, et al. American Society of Hematology 2019 guidelines for management of venous thromboembolism: prevention of venous thromboembolism in surgical hospitalized patients. *Blood Adv*. 2019;3(23):3898-3944. doi:10.1182/bloodadvances.2019000975
3. CRISTAL Study Group, Sidhu VS, Kelly TL, et al. Effect of Aspirin vs Enoxaparin on Symptomatic Venous Thromboembolism in Patients Undergoing Hip or Knee Arthroplasty: The CRISTAL Randomized Trial. *JAMA*. 2022;328(8):719-727. doi:10.1001/jama.2022.13416
4. Anderson DR, Dunbar M, Murnaghan J, et al. Aspirin or Rivaroxaban for VTE Prophylaxis after Hip or Knee Arthroplasty. *N Engl J Med*. 2018;378(8):699-707. doi:10.1056/NEJMoa1712746
5. Fagarasanu A, Alotaibi GS, Hrimiuc R, Lee AYY, Wu C. Role of Extended Thromboprophylaxis After Abdominal and Pelvic Surgery in Cancer Patients: A Systematic Review and Meta-Analysis. *Ann Surg Oncol*. 2016;23(5):1422-1430. doi:10.1245/s10434-016-5127-1
6. Aminian A, Vosburg RW, Altieri MS, Hinojosa MW, Khorgami Z, American Society for Metabolic and Bariatric Surgery Clinical Issues Committee. The American Society for Metabolic and Bariatric Surgery (ASMBS) updated position statement on perioperative venous thromboembolism prophylaxis in bariatric surgery. *Surg Obes Relat Dis Off J Am Soc Bariatr Surg*. 2022;18(2):165-174. doi:10.1016/j.soard.2021.10.023
7. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336(7650):924-926. doi:10.1136/bmj.39489.470347.AD
8. Falck-Ytter Y, Francis CW, Johanson NA, et al. Prevention of VTE in orthopedic surgery patients: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*. 2012;141(2 Suppl):e278S-e325S. doi:10.1378/chest.11-2404
9. Caprini JA. Risk assessment as a guide to thrombosis prophylaxis. *Curr Opin Pulm Med*. 2010;16(5):448-452. doi:10.1097/MCP.0b013e32833c3d3e

10. Lindblad B, Eriksson A, Bergqvist D. Autopsy-verified pulmonary embolism in a surgical department: analysis of the period from 1951 to 1988. *Br J Surg*. 1991;78(7):849-852. doi:10.1002/bjs.1800780725
11. Stein PD, Henry JW. Prevalence of acute pulmonary embolism among patients in a general hospital and at autopsy. *Chest*. 1995;108(4):978-981. doi:10.1378/chest.108.4.978
12. White RH, Zhou H, Romano PS. Incidence of symptomatic venous thromboembolism after different elective or urgent surgical procedures. *Thromb Haemost*. 2003;90(3):446-455. doi:10.1160/TH03-03-0152
13. Buesing KL, Mullapudi B, Flowers KA. Deep venous thrombosis and venous thromboembolism prophylaxis. *Surg Clin North Am*. 2015;95(2):285-300. doi:10.1016/j.suc.2014.11.005
14. Gordon RJ, Lombard FW. Perioperative Venous Thromboembolism: A Review. *Anesth Analg*. 2017;125(2):403-412. doi:10.1213/ANE.0000000000002183
15. Trivedi NN, Abola MV, Kim CY, Sivasundaram L, Smith EJ, Ochenjele G. The Incremental Cost of Inpatient Venous Thromboembolism After Hip Fracture Surgery. *J Orthop Trauma*. 2020;34(4):169-173. doi:10.1097/BOT.0000000000001675
16. Ollendorf DA, Vera-Llonch M, Oster G. Cost of venous thromboembolism following major orthopedic surgery in hospitalized patients. *Am J Health-Syst Pharm AJHP Off J Am Soc Health-Syst Pharm*. 2002;59(18):1750-1754. doi:10.1093/ajhp/59.18.1750
17. Office of the Surgeon General (US), National Heart, Lung, and Blood Institute (US). *The Surgeon General's Call to Action to Prevent Deep Vein Thrombosis and Pulmonary Embolism*. Office of the Surgeon General (US); 2008. Accessed October 15, 2023. <http://www.ncbi.nlm.nih.gov/books/NBK44178/>
18. Streiff MB, Haut ER. The CMS ruling on venous thromboembolism after total knee or hip arthroplasty: weighing risks and benefits. *JAMA*. 2009;301(10):1063-1065. doi:10.1001/jama.301.10.1063
19. Roberts VC, Sabri S, Beeley AH, Cotton LT. The effect of intermittently applied external pressure on the haemodynamics of the lower limb in man. *Br J Surg*. 1972;59(3):223-226. doi:10.1002/bjs.1800590319
20. Comerota AJ, Chouhan V, Harada RN, et al. The fibrinolytic effects of intermittent pneumatic compression: mechanism of enhanced fibrinolysis. *Ann Surg*. 1997;226(3):306-313; discussion 313-314. doi:10.1097/0000658-199709000-00010
21. Kakkos SK, Caprini JA, Geroulakos G, Nicolaidis AN, Stansby GP, Reddy DJ. Combined intermittent pneumatic leg compression and pharmacological prophylaxis for prevention of venous thromboembolism in high-risk patients. *Cochrane Database Syst Rev*. 2008;(4):CD005258. doi:10.1002/14651858.CD005258.pub2

22. Eppsteiner RW, Shin JJ, Johnson J, van Dam RM. Mechanical compression versus subcutaneous heparin therapy in postoperative and posttrauma patients: a systematic review and meta-analysis. *World J Surg*. 2010;34(1):10-19. doi:10.1007/s00268-009-0284-z
23. Ho KM, Tan JA. Stratified meta-analysis of intermittent pneumatic compression of the lower limbs to prevent venous thromboembolism in hospitalized patients. *Circulation*. 2013;128(9):1003-1020. doi:10.1161/CIRCULATIONAHA.113.002690
24. Camporese G, Bernardi E, Prandoni P, et al. Low-molecular-weight heparin versus compression stockings for thromboprophylaxis after knee arthroscopy: a randomized trial. *Ann Intern Med*. 2008;149(2):73-82. doi:10.7326/0003-4819-149-2-200807150-00003
25. Morris RJ, Woodcock JP. Intermittent pneumatic compression or graduated compression stockings for deep vein thrombosis prophylaxis? A systematic review of direct clinical comparisons. *Ann Surg*. 2010;251(3):393-396. doi:10.1097/SLA.0b013e3181b5d61c
26. Arabi YM, Khedr M, Dara SI, et al. Use of intermittent pneumatic compression and not graduated compression stockings is associated with lower incident VTE in critically ill patients: a multiple propensity scores adjusted analysis. *Chest*. 2013;144(1):152-159. doi:10.1378/chest.12-2028
27. Kaw R, Pasupuleti V, Wayne Overby D, et al. Inferior vena cava filters and postoperative outcomes in patients undergoing bariatric surgery: a meta-analysis. *Surg Obes Relat Dis Off J Am Soc Bariatr Surg*. 2014;10(4):725-733. doi:10.1016/j.soard.2014.04.008
28. Ikesaka R, Kaur B, Crowther M, Rajasekhar A. Efficacy and safety of pre-operative insertion of inferior vena cava filter in patients undergoing bariatric surgery: a systematic review. *J Thromb Thrombolysis*. 2022;54(3):502-523. doi:10.1007/s11239-022-02689-3
29. Ho KM, Rao S, Honeybul S, et al. A Multicenter Trial of Vena Cava Filters in Severely Injured Patients. *N Engl J Med*. 2019;381(4):328-337. doi:10.1056/NEJMoa1806515
30. Hemmila MR, Osborne NH, Henke PK, et al. Prophylactic Inferior Vena Cava Filter Placement Does Not Result in a Survival Benefit for Trauma Patients. *Ann Surg*. 2015;262(4):577-585. doi:10.1097/SLA.0000000000001434
31. Angel LF, Tapson V, Galgon RE, Restrepo MI, Kaufman J. Systematic review of the use of retrievable inferior vena cava filters. *J Vasc Interv Radiol JVIR*. 2011;22(11):1522-1530.e3. doi:10.1016/j.jvir.2011.08.024
32. Kaufman JA, Barnes GD, Chaer RA, et al. Society of Interventional Radiology Clinical Practice Guideline for Inferior Vena Cava Filters in the Treatment of Patients with Venous Thromboembolic Disease: Developed in collaboration with the American College of Cardiology, American College of Chest Physicians, American College of Surgeons Committee on Trauma, American Heart Association, Society for Vascular Surgery, and Society for

Vascular Medicine. *J Vasc Interv Radiol JVIR*. 2020;31(10):1529-1544.  
doi:10.1016/j.jvir.2020.06.014

33. Safety Communications > Removing Retrievable Inferior Vena Cava Filters: FDA Safety Communication. Accessed October 15, 2023. <http://wayback.archive-it.org/7993/20170722215731/https://www.fda.gov/MedicalDevices/Safety/AlertsandNotices/ucm396377.htm>
34. Prevention of pulmonary embolism and deep vein thrombosis with low dose aspirin: Pulmonary Embolism Prevention (PEP) trial. *Lancet Lond Engl*. 2000;355(9212):1295-1302.
35. Matharu GS, Kunutsor SK, Judge A, Blom AW, Whitehouse MR. Clinical Effectiveness and Safety of Aspirin for Venous Thromboembolism Prophylaxis After Total Hip and Knee Replacement: A Systematic Review and Meta-analysis of Randomized Clinical Trials. *JAMA Intern Med*. 2020;180(3):376-384. doi:10.1001/jamainternmed.2019.6108
36. Anderson DR, Dunbar MJ, Bohm ER, et al. Aspirin versus low-molecular-weight heparin for extended venous thromboembolism prophylaxis after total hip arthroplasty: a randomized trial. *Ann Intern Med*. 2013;158(11):800-806. doi:10.7326/0003-4819-158-11-201306040-00004
37. Major Extremity Trauma Research Consortium (METRC), O'Toole RV, Stein DM, et al. Aspirin or Low-Molecular-Weight Heparin for Thromboprophylaxis after a Fracture. *N Engl J Med*. 2023;388(3):203-213. doi:10.1056/NEJMoa2205973
38. Drescher FS, Sirovich BE, Lee A, Morrison DH, Chiang WH, Larson RJ. Aspirin versus anticoagulation for prevention of venous thromboembolism major lower extremity orthopedic surgery: a systematic review and meta-analysis. *J Hosp Med*. 2014;9(9):579-585. doi:10.1002/jhm.2224
39. Pellegrini VD. *Comparative Effectiveness of Pulmonary Embolism Prevention After Hip and Knee Replacement: Balancing Safety and Effectiveness*. clinicaltrials.gov; 2022. Accessed December 31, 2022. <https://clinicaltrials.gov/study/NCT02810704>
40. Shivakumar S. *Extended Venous Thromboembolism Prophylaxis Comparing Rivaroxaban and Aspirin to Aspirin Alone Following Total Hip and Knee Arthroplasty (EPCATIII)*. clinicaltrials.gov; 2022. Accessed December 31, 2022. <https://clinicaltrials.gov/study/NCT04075240>
41. Turpie AG, Levine MN, Hirsh J, et al. A randomized controlled trial of a low-molecular-weight heparin (enoxaparin) to prevent deep-vein thrombosis in patients undergoing elective hip surgery. *N Engl J Med*. 1986;315(15):925-929. doi:10.1056/NEJM198610093151503

42. Leclerc JR, Geerts WH, Desjardins L, et al. Prevention of deep vein thrombosis after major knee surgery--a randomized, double-blind trial comparing a low molecular weight heparin fragment (enoxaparin) to placebo. *Thromb Haemost*. 1992;67(4):417-423.
43. Fuji T, Ochi T, Niwa S, Fujita S. Prevention of postoperative venous thromboembolism in Japanese patients undergoing total hip or knee arthroplasty: two randomized, double-blind, placebo-controlled studies with three dosage regimens of enoxaparin. *J Orthop Sci Off J Jpn Orthop Assoc*. 2008;13(5):442-451. doi:10.1007/s00776-008-1264-0
44. Hill J, Treasure T, National Clinical Guideline Centre for Acute and Chronic Conditions. Reducing the risk of venous thromboembolism in patients admitted to hospital: summary of NICE guidance. *BMJ*. 2010;340:c95. doi:10.1136/bmj.c95
45. Mismetti P, Laporte S, Zufferey P, Epinat M, Decousus H, Cucherat M. Prevention of venous thromboembolism in orthopedic surgery with vitamin K antagonists: a meta-analysis. *J Thromb Haemost JTH*. 2004;2(7):1058-1070. doi:10.1111/j.1538-7836.2004.00757.x
46. Kapoor A, Ellis A, Shaffer N, et al. Comparative effectiveness of venous thromboembolism prophylaxis options for the patient undergoing total hip and knee replacement: a network meta-analysis. *J Thromb Haemost JTH*. 2017;15(2):284-294. doi:10.1111/jth.13566
47. Lassen MR, Raskob GE, Gallus A, Pineo G, Chen D, Portman RJ. Apixaban or enoxaparin for thromboprophylaxis after knee replacement. *N Engl J Med*. 2009;361(6):594-604. doi:10.1056/NEJMoa0810773
48. Kakkar AK, Brenner B, Dahl OE, et al. Extended duration rivaroxaban versus short-term enoxaparin for the prevention of venous thromboembolism after total hip arthroplasty: a double-blind, randomised controlled trial. *Lancet Lond Engl*. 2008;372(9632):31-39. doi:10.1016/S0140-6736(08)60880-6
49. Eriksson BI, Bauer KA, Lassen MR, Turpie AG, Steering Committee of the Pentasaccharide in Hip-Fracture Surgery Study. Fondaparinux compared with enoxaparin for the prevention of venous thromboembolism after hip-fracture surgery. *N Engl J Med*. 2001;345(18):1298-1304. doi:10.1056/NEJMoa011100
50. Lassen MR, Gallus A, Raskob GE, et al. Apixaban versus enoxaparin for thromboprophylaxis after hip replacement. *N Engl J Med*. 2010;363(26):2487-2498. doi:10.1056/NEJMoa1006885
51. Eriksson BI, Dahl OE, Rosencher N, et al. Oral dabigatran etexilate vs. subcutaneous enoxaparin for the prevention of venous thromboembolism after total knee replacement: the RE-MODEL randomized trial. *J Thromb Haemost JTH*. 2007;5(11):2178-2185. doi:10.1111/j.1538-7836.2007.02748.x
52. Eriksson BI, Borris LC, Friedman RJ, et al. Rivaroxaban versus enoxaparin for thromboprophylaxis after hip arthroplasty. *N Engl J Med*. 2008;358(26):2765-2775. doi:10.1056/NEJMoa0800374

53. Hull RD, Pineo GF, Stein PD, et al. Extended out-of-hospital low-molecular-weight heparin prophylaxis against deep venous thrombosis in patients after elective hip arthroplasty: a systematic review. *Ann Intern Med.* 2001;135(10):858-869. doi:10.7326/0003-4819-135-10-200111200-00006
54. Petersen PB, Kehlet H, Jørgensen CC, Lundbeck Foundation Centre for Fast-track Hip and Knee Replacement Collaborative Group. Safety of In-Hospital Only Thromboprophylaxis after Fast-Track Total Hip and Knee Arthroplasty: A Prospective Follow-Up Study in 17,582 Procedures. *Thromb Haemost.* 2018;118(12):2152-2161. doi:10.1055/s-0038-1675641
55. Dahl OE, Gudmundsen TE, Pripp AH, Aanesen JJ. Clinical venous thromboembolism following joint surgery: effect of extended thromboprophylaxis on its annual frequency and postoperative pattern over 22 years. *Clin Appl Thromb Off J Int Acad Clin Appl Thromb.* 2014;20(2):117-123. doi:10.1177/1076029613499820
56. Partridge T, Jameson S, Baker P, Deehan D, Mason J, Reed MR. Ten-Year Trends in Medical Complications Following 540,623 Primary Total Hip Replacements from a National Database. *J Bone Joint Surg Am.* 2018;100(5):360-367. doi:10.2106/JBJS.16.01198
57. Selby R, Geerts WH, Kreder HJ, Crowther MA, Kaus L, Sealey F. Symptomatic venous thromboembolism uncommon without thromboprophylaxis after isolated lower-limb fracture: the knee-to-ankle fracture (KAF) cohort study. *J Bone Joint Surg Am.* 2014;96(10):e83. doi:10.2106/JBJS.M.00236
58. Wahlsten LR, Eckardt H, Lyngbæk S, et al. Symptomatic venous thromboembolism following fractures distal to the knee: a nationwide Danish cohort study. *J Bone Joint Surg Am.* 2015;97(6):470-477. doi:10.2106/JBJS.N.00307
59. van Adrichem RA, Nemeth B, Algra A, et al. Thromboprophylaxis after Knee Arthroscopy and Lower-Leg Casting. *N Engl J Med.* 2017;376(6):515-525. doi:10.1056/NEJMoa1613303
60. Nemeth B, Timp JF, van Hylckama Vlieg A, Rosendaal FR, Cannegieter SC. High risk of recurrent venous thrombosis in patients with lower-leg cast immobilization. *J Thromb Haemost JTH.* 2018;16(11):2218-2222. doi:10.1111/jth.14278
61. Bahl V, Hu HM, Henke PK, Wakefield TW, Campbell DA, Caprini JA. A validation study of a retrospective venous thromboembolism risk scoring method. *Ann Surg.* 2010;251(2):344-350. doi:10.1097/SLA.0b013e3181b7fca6
62. Anderson FA, Spencer FA. Risk Factors for Venous Thromboembolism. *Circulation.* 2003;107(23\_suppl\_1):I-9. doi:10.1161/01.CIR.0000078469.07362.E6
63. Caprini JA, Arcelus JJ, Hasty JH, Tamhane AC, Fabrega F. Clinical assessment of venous thromboembolic risk in surgical patients. *Semin Thromb Hemost.* 1991;17 Suppl 3:304-312.
64. Rogers SO, Kilaru RK, Hosokawa P, Henderson WG, Zinner MJ, Khuri SF. Multivariable predictors of postoperative venous thromboembolic events after general and vascular

surgery: results from the patient safety in surgery study. *J Am Coll Surg*. 2007;204(6):1211-1221. doi:10.1016/j.jamcollsurg.2007.02.072

65. Mulder FI, Horváth-Puhó E, van Es N, et al. Venous thromboembolism in cancer patients: a population-based cohort study. *Blood*. 2021;137(14):1959-1969. doi:10.1182/blood.2020007338
66. Lyman GH, Carrier M, Ay C, et al. American Society of Hematology 2021 guidelines for management of venous thromboembolism: prevention and treatment in patients with cancer. *Blood Adv*. 2021;5(4):927-974. doi:10.1182/bloodadvances.2020003442
67. Aminian A, Andalib A, Khorgami Z, et al. Who Should Get Extended Thromboprophylaxis After Bariatric Surgery?: A Risk Assessment Tool to Guide Indications for Post-discharge Pharmacoprophylaxis. *Ann Surg*. 2017;265(1):143-150. doi:10.1097/SLA.0000000000001686
68. Thereaux J, Lesuffleur T, Czernichow S, et al. To What Extent Does Posthospital Discharge Chemoprophylaxis Prevent Venous Thromboembolism After Bariatric Surgery?: Results From a Nationwide Cohort of More Than 110,000 Patients. *Ann Surg*. 2018;267(4):727-733. doi:10.1097/SLA.0000000000002285
69. Kröll D, Nett PC, Rommers N, et al. Efficacy and Safety of Rivaroxaban for Postoperative Thromboprophylaxis in Patients After Bariatric Surgery: A Randomized Clinical Trial. *JAMA Netw Open*. 2023;6(5):e2315241. doi:10.1001/jamanetworkopen.2023.15241
70. Hemmila MR, Osborne NH, Henke PK, Kepros JP, Patel SG, Cain-Nielsen AH, Birkmeyer NJ. Inferior Vena Cava Filter Placement Does Not Result in a Survival Benefit for Trauma Patients. *Ann Surg*. 2015 Oct;262(4):577-85. doi: 10.1097/SLA.0000000000001434. PMID: 26366537.
71. Sarosiek S, Rybin D, Weinberg J, Burke PA, Kasotakis G, Sloan JM. Association Between Inferior Vena Cava Filter Insertion in Trauma Patients and In-Hospital and Overall Mortality. *JAMA Surg*. 2017 Jan 1;152(1):75-81. doi: 10.1001/jamasurg.2016.3091. PMID: 27682367.